


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

Scholar [All articles](#) - [Recent articles](#) Results 11 - 20 of about 2,000 for spiked arabidopsis. (0.11 s)

[PDF] [Empirical evaluation of methodologies for microarray data analysis - all 3 versions »](#)

L Qin, K Kerr - Nucleic Acids Res, 2004 - ima.umn.edu

... Arabidopsis genes spiked in • 4 genes spiked in at 1:1 ratio ... l: 4 Arabidopsis spike-ins at differential ratios 48 spots for each spike-in ...

[Cited by 2](#) - [Related Articles](#) - [View as HTML](#) - [Web Search](#)

[Accumulation of 6-deoxocathasterone and 6-deoxocastasterone in Arabidopsis, pea and tomato is ... - all 5 versions »](#)

T Nomura, T Sato, GJ Bishop, Y Kamiya, S Takatsuto ... - Phytochemistry, 2001 - Elsevier

... The MeOH extract of the harvested materials was spiked with 2 H 6-labelled BRs as ... CHCl 3 containing 0, 8, and 10% MeOH in the case of Arabidopsis, with CHCl 3 ...

[Cited by 34](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[... indole-3-acetic acid biosynthesis in the wild-type and a tryptophan auxotroph mutant of Arabidopsis ... - all 3 versions »](#)

A Müller, EW Weiler - Planta, 2000 - Springer

... in the wild-type and a tryptophan auxotroph mutant of Arabidopsis thaliana ... Plant material and growth conditions Arabidopsis thaliana (L.) Heynh. ...

[Cited by 22](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Analysis of the defence phosphoproteome of Arabidopsis thaliana using differential mass tagging - all 3 versions »](#)

AM Jones, MH Bennett, JW Mansfield, M Grant - Proteomics, 2006 - doi.wiley.com

... [17] who used three yeast strains and spiked extracts with ... peptides during the early stages of the response of the model plant Arabidopsis thaliana to ...

[Cited by 22](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Hyperaccumulation of Cadmium and Zinc in Thlaspi caerulescens and Arabidopsis halleri at the Leaf ... - all 6 versions »](#)

C Cosio, E Martinoia, C Keller - Plant Physiology, 2004 - pubmedcentral.nih.gov

... al., 2000), T. caerulescens ecotype "Ganges" (southern France) and Arabidopsis halleri (northern ... was initiated by the addition of ZnCl 2 spiked with 18.5 ...

[Cited by 31](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[... and absolute quantitative shotgun proteomics: targeting low-abundance proteins in Arabidopsis ... - all 6 versions »](#)

S Wienkoop, W Weckwerth - Journal of Experimental Botany, 2006 - Soc Experiment Biol

... 4). For peak integration, three replicates of 20 µg digested Arabidopsis protein extract with and without 500 fmol spiked standard peptide each were analysed ...

[Cited by 8](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Identification of Arabidopsis salt and osmotic stress responsive proteins using two-dimensional ... - all 3 versions »](#)

BK Ndimba, S Chivasa, WJ Simon, AR Slabas - Proteomics, 2005 - doi.wiley.com

... Arabidopsis cell suspension cultures were simul- taneously spiked with 35 S-methionine/ 35 S cysteine and either water (Cn), 200m M NaCl (S) or 400m M sorbitol ...

[Cited by 15](#) - [Related Articles](#) - [Web Search](#)